8000124

THE UNITED SEATES OF AVIERA CA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Dairyland Seed Co., Inc.

Withereas, there has been presented to the

Досточной сов чубане в сан цана с

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic ed of the variety in a public repository as provided by LAW, the right to exothers from selling the variety, or offering it for sale, or reproducing it, or therefrom, to the extent provided by the Plant Variety Protection Act.

In therefrom, to the extent provided by the Plant Variety Protection Act.

In the States seed of this variety (1) shall be sold by variety name only as certified seed and (2) shall conform to the number of generations the owner of the rights. (84 stat. 1542, as amended, 7 u.s.c. 2321 et seq.)

RED CLOVER

'Ruby'

In Testimony Winexcot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington

at the City of Washington
this 15th day of April in
the year of our Lord one thousand nine
hundred and eighty-two.

Attest:

Kennett H.E.

'Acting Commissione

Diant Variety Protection Office G . Or . .

Agricultural Marketing Service

Heretary of Agriculture

AP	UNITED STATES DEPARTME AGRICULTURAL MARY LIVESTOCK, POULTRY, GRA PLICATION FOR PLANT VARIE	CETING SERVICE VIN & SEED DIVISION		be issued unless a co	OMB NO. 40-R38 ant variety protection m impleted application for	22 ay
	TRUCTIONS: See Reverse. TEMPORARY DESIGNATION OF	1b. VARIETY NAM	E			
	JOY	RUBY	• :	PV NUMBER		_
2.	KIND NAME	3. GENUS AND SPE	CIES NAME	FILING DATE		<u>—</u>
	RED CLOVER	TRIFOLIUM	PRATENSE	5/16/80	P.1	À,
4.	FAMILY NAME (BOTANICAL)	5. DATE OF DETER	NOITANIME	s 500.00	5/16/80	
	LEGUMINOSAE	JANUARY,	19 78	\$ 250.00	12/11/81	
6.	NAME OF APPLICANT(S)	7. ADDRESS (Street Code)	and No. or R.F.D. No.,	City, State, and ZIP		
	DAIRYLAND SEED CO., INC.	P.O. BOX WEST BEND		4		
9.	IF THE NAMED APPLICANT IS NOT A PE	RSON, FORM OF		FOR OFFICIAL USE ONLY PV NUMBER 8000124 FILING DATE 5/16/80 FILING 6/16/80 FIL		
	ORGANIZATION: (Corporation, partnersh CORPORATION	ip, association, etc.)	WISCONSIN	PORATION	OMB NO. 40-R3822 Occrificate for plant variety protection may issued unless a completed application form is been received (5 U.S.C. 552). FOR OFFICIAL USE ONLY IUMBER 8000124 NG DATE	
12.	NAME AND MAILING ADDRESS OF APP ALL PAPERS:	LICANT REPRESENTA	ATIVE(S), IF ANY, TO S	SERVE IN THIS APPLIC	ATION AND RECEIVE	_
М	R. THOMAS G. STRACHOTA, DAT	IRYLAND SEED CO	O., INC. P.O.	BOX 958 WEST	BEND, WI 5309	5
13.	CHECK BOX BELOW FOR EACH ATTACH	MENT SUBMITTED:				-
	13B. Exhibit B, Novelty Statem	ent.				
	13D. Exhibit D, Additional Desc	and the second second	No	Plant Variety Protect	ion Office.)	
14a,	DOES THE APPLICANT(S) SPECIFY THAT SEED? (See Section 83(a). (If "Yes," answer	T SEED OF THIS VARI er 14B and 14C below.)			A CLASS OF CERTIFI	ED
14b.	DOES THE APPLICANT(S) SPECIFY THAT LIMITED AS TO NUMBER OF GENERATI	T THIS VARIETY BE	14c. IF "YES," TO 14I	B, HOW MANY GENERA	ATIONS OF PRODUC-	A.M. P.M. A.M. P.M. 80 81 NE AREA D NUMBER 8-0163 FINCOR- DN 1963 RECEIVE 53095 Act.) CERTIFIED COUC- ED Yes, "give
	X YES NO		X FOUNDATION		X CERTIFIED	
15a,	DID THE APPLICANT(S) FILE FOR PROT name of countries and dates.)	ECTION OF THIS VAF	RIETY IN OTHER COUN	NTRIES? YES	X NO (If "Yes," giv	ie
				HER WITH BUILDING		
15b.	HAVE RIGHTS BEEN GRANTED THIS VA and dates.)	RIETY IN OTHER CO	UNTRIES? TYES	X NO (If "Yes,"	give name of countries	
			The state of the s			
16.	DOES THE APPLICANT(S) AGREE TO THE JOURNAL?	E PUBLICATION OF H	IS/HER (THEIR) NAME	(S) AND ADDRESS IN	THE OFFICIAL	-
17.		sample of basic seed	l of this variety will b	e furnished with the a	pplication and will be	-
	The undersigned applicant(s) is (are) th	e owner(s) of this se	xually reproduced nov	vel plant variety, and	believe(s) that the provisions of Section	1
	Applicant(s) is (are) informed that false	e representation here	in can jeopardize prot	ection and result in p	enalties.	
	4-15-80		Momas	Strack	tota	_
	(DATE)		(S	SIGNATURE OF APPLIC	CANT)	_

INSTRUCTIONS

GENERAL: Send an original copy of the application and exhibits, at least 2,500 viable seeds, and \$500 fee (\$250 filing fee and \$250 examination fee) to U.S. Dept. of Agriculture, Agricultural Marketing Service, Livestock, Poultry, Grain and Seed Division, Plant Variety Protection Office, National Agricultural Library Building, Beltsville, Maryland 20705. (See section 180.175 of the Regulations and Rules of Practice.) Retain one copy for your files. All items on the face of the form are self-explanatory unless noted below.

ITEM

- Give the date the applicant determined that he had a new variety based on (1) the definition in section 41(a) of the Act and (2) the date a decision was made to increase the seed.
- Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method; (2) the details of subsequent stages of selection and multiplication; (3) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified and (4) evidence of uniformity and stability.
- Give a summary statement of the variety's novelty. Clearly state how this novel variety may be distinguished from all other varieties in the same crop. If the new variety most closely resembles one or a group of related varieties:

 (1) identify these varieties and state all differences objectively; (2) attach statistical data for characters expressed numerically and demonstrate that these differences are significant; and (3) submit, if helpful, seed and plant specimens or photographs of seed and plant comparisons clearly indicating novelty.
- Fill in the Exhibit C, Objective Description form, for all characteristics for which you have adequate data.
- Describe any additional characteristics that are not described, or whose description cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the description of characteristics that are difficult to describe, such as, plant habit, plant color, disease resistance, etc.
- If "YES" is specified (seed of this variety be sold by variety name only as a class of certified seed) the applicant may NOT reverse his affirmative decision after the variety has either been sold and so labeled, his decision published, or the certificate has been issued. However, if the applicant specified "NO," he may change his choice. (See section 180.16 of the Regulations and Rules of Practice.)
- See section 42 of the Plant Variety Protection Act and section 180.7 of the Regulations and Rules of Practice.

Ruby was developed under the experimental name Joy by the Teweles Seed Company at its Clinton, Wisconsin Research Station and at its Sloughhouse, California Research Station. In November 1977 Dairyland Seed Co., Inc. purchased the Teweles Research facilities and acquired rights to this red clover variety. Dairyland then changed the name to Ruby and released the variety. Parental sources were selected from Arlington and Kenstar based on progency tests for forage yield, seed yield, forage productivity persistance, winter hardiness, and disease resistance. Forage and disease research was conducted at the Clinton Research Station. Seed set research was conducted at the Sloughhouse Research Station. Ruby has been observed over generations and is a stable variety that does not change in its essential and distinctive characters. Ruby is as uniform as other varieties of red clover. No variants have been observed outside the range normal for a cross-pollinated variety. A sufficient quantity of breeder and foundation seed has been increased for the life of the variety. This seed is being maintained under climate controlled conditions. There are three classes of seed increase: breeder seed, foundation seed, and certified seed.

RUBY RED CLOVER

EXHIBIT B

NOVELTY STATEMENT

Ruby Red Clover is most similar to Kenstar Red Clover. However, Ruby has both northern and southern anthracnose resistance, better powdery mildew resistance, and more winter hardy than Kenstar. Kenstar is resistant to southern anthracnose but susceptible to northern anthracnose whereas Arlington is resistant to northern anthracnose but susceptible to southern anthracnose. This additional disease resistance and winter hardiness makes Ruby have a much wider range of geographic adaptation which would be considered a novel trait of important agricultural value.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE GRAIN AND SEED DIVISION BELTSVILLE, MARYLAND 20705

(Red Clover)

OBJECTIVE DESCRIPTION OF VARIETY

RED CLOVER (Trifolium Pratense)

NAME OF APPLICANT(S) DAIRYLAND SEED COMPANY, INC.	VARIETY NAME OR TEMPORARY DESIGNATION RUBY
ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code)	FOR OFFICIAL USE ONLY
P. O. BOX 958	PVPO NUMBER
WEST BEND, WISCONSIN 53095	8000124
Place the appropriate number that describes the varietal character of this variety in the boxes when number is 99). In comparisons to standard varieties, the value 0 0 should only symbol indicates a decimal point. Characteristics described, including numerical measure the variety. Measured data should be for SPACED PLANTS. Any recognized color fan, e.g. mine plant colors; designate system used: Ranges of values are valuable and may be included with additional description elsewhere in the NOTE: For single plant data a minimum of 100 plants is suggested. 1. TYPE:	be used to indicate that the varieties are equal. The ements, should represent those which are TYPICAL for Royal Horticultural Colour Chart, may be used to dete
1 1 = DOUBLE CUT (medium) 2 = SINGLE CUT (mammoth)	3 = OTHER (Specify)
2. PLOIDY:	
1 = DIPLOID 2 = TETRAPLOID	3 = OTHER (Specify)
3. PRODUCTIVE PERSISTENCE (Usual duration of planting):	
2 1 = ANNUAL 2 = BIENNIAL 3 = SHORT	LIVED PERENNIAL (3 - 4 Years)
4. ADAPTATION: (e.g., 0 2 3 = northcentral and southcentral)	
0 0 2 1 = NORTHEAST 2 = NORTHCENTRAL	3 = SOUTHCENTRAL
4 = SOUTHEAST 5 = WEST	6 = OTHER (Specify)
STANDARD VARIETIES	
1 = KENSTAR 2 = ARLINGTON 3 = PENNSCOTT	4 = TENSAS 5 = ALTASWEDE
5. MATURITY: 0 11/24/8/ 0 79 5 % PLANTS FLOWERING IN SEEDLING YEAR	
Beginning of spring growth:	
0 2 DAYS EARLIER THAN 2 STANDAR	D VARIETY
DAYS LATER THAN STANDAR	D VARIETY
Time of flowering (50% of plants in bloom): (from spring growth in non-se	edling year)
0 3 DAYS EARLIER THAN	D VARIETY
DAYS LATER THAN STANDAR	D VARIETY
6. PLANT HEIGHT (from soil level to top of flowering head at 50% flowering)	
* 6 7 CM. TALL CM. SHOR	FER THAN STANDARD VARIETY
1 6 CM. TALLE	R THAN 1 STANDARD VARIETY
* Shorter than the total length of stem because of prostration at the base of the stem.	

<u> F</u>	ORM GR-470-4	2				
-	7. FLOWER	ING STEM (from first none	contracted internode	e, longer than 0.	5 cm., to tip of flowering he	ad):
	2	NO. FLOWERING STEMS	PER CROWN			
	8	NO. INTERNODES				
	7 3	CM. LENGTH OF STEM	·			
	Hairiness	Give percentage of plants	with each type of s	urface (Total =	100%)	
		% HAIRS PROJECTING U	PWARD			
	> 9 9	% HAIRS PROJECTING D	OWNWARD OR AT R	IGHT ANGLES		•
		% GLABROUS (FEWER TI	•			
e		ve percentage of plants wit he horizontal (soil level) at		. Stem habit sh	ould be determined by the a	ingle of lowest
1124/	¥ 9 9	% PROSTRATE (0 - 30°) (1/2)	1 % SEMI-PROST (30 - 45°)		0 % SEMI-ERECT (45 - 60°)	0 % ERECT (60 - 90°)
	B. LEAF (Cei	ntral leaflet at 3rd node bel	ow flowering head):			•
	2 0	MM WIDTH		IARROWER THA		ARD VARIETY ARD VARIETY
4	4 1	MM LENGTH		HORTER THAN		ARD VARIETY
	Color:					
	2	1 = LIGHT GREEN (Altaswede)	2 = MEDIUM GI	REEN)	3 = DARK GREEN (Hungaropoli)	4 = BLUE GREEN
		Leaf Marking (at 50% flo data. Diagram illustrates 2D = INCOMPLETE 3	terms: 1 = APICA	ategories below L 2A = FULL	allow for increasingly detail 2B = EXTENDED 2C = APICAL CENTRA	led description of the same DELTA BASAL
		f Mark: Of total plants, gi (Total = 100%)	ve percentage marke	ed and	0000	
	2 2	% ABSENT 7	8 % MARKED		1 2A 2B 2	C 2D 3
	Position of	Mark: Of total plants, giv	e percentage with le	af mark in each	position (Total = % marked	, above)
[8	% APICAL		ITRAL	0 % BASA	
	Shape of N	lark: Of total plants, give	percentage with cen	tral leaf marks h	aving each shape (Total = %	marked, above)
	0	% FULL	0 % EXTENDED	6 2	% DELTA 8	% INCOMPLETE
g		OLOR (Determine color o eferenced to the Munsell C		rets): Give per	centage of plants with each o	color (Total = 100%).
		% WHITE		2 3	% LIGHT PINK (5RP 8/4)	
	7 7	% MEDIUM PINK (5RP 7/6)			% DARK PINK (5RP 6/8)	•
ſ		% RED (5RP 5/10)			% OTHER (Specify)	

FORM GR-470-42	***************************************		· · · · · · · · · · · · · · · · · · ·	Page 3 o
10. SEED COLOR: Maxim plants with each seed co	num color development in unstored	d, mature seed	(at beginning of ca	lyx browning). Give percentage of
plants with duen seed to	100 // 100 //		•	
4 4 % YELLOW	v .	1 7	% YELLOW WITH	SOME PURPLE
1 3 % PURPLE	[2 6	% PURPLE WITH S	OME YELLOW
% OTHER (Specify)		_ (attach explanation)	
11. DISEASE AND INSECT	RESISTANCE (0 = not tested, 1	= susceptible	, and 2 = resistant).	If variety is claimed to be resistant
or to show intermediate	reaction, substantiating test score location of test, and range and di	s should be at	tached clearly ident	ifying disease, application variety,
	i location of test, and range and of	rection of test	scores.	
A. DISEASES:				•
CROWN RO	T (Sclerotina trifoliorum)		ROOT ROT (Fusario	um spp.)
2 NORTHERN	N ANTHRACNOSE (Kabatiella caulivoi	ral	SUMMER BLACK S	TEM (Cercospora zebrina)
SOUTHERN	ANTHRACNOSE (Colletotrichum tri	folii)	BLACK STEM (Phor	na trifolii)
TARGET SP	OT (Stemphylium sarcinaeformae)	2	POWDERY MILDEN	N (Erysiphe polygoni)
PEPPER SPO	T (Leptosphaeralina trifolii)		BLACK PATCH (Rh	izoctonia leguminicola)
RED CLOVE	R VEIN MOSIAC VIRUS		BEAN YELLOW MO	SIAC VIRUS
NEMATODE	(Specify)		OTHER (Specify)	-
B. INSECTS:	•			
CLOVER RO	OT BORER (Hylastinus obsurus)		CLOVER ROOT CUI	RCULIO (Sitona hispidula)
SWEETCLOV	ER WEEVIL (Sitona cylindricollis)		CLOVER SEED CHA	LCID (Bruchophagus platyptera)
LESSER CLO	VER LEAF WEEVIL		POTATO LEAFHOPE	PER (Empoasca fabae)
(Hypera nigrir	*		MEADOW SPITTLES	SUG (Philaenus spumarius)
YELLOW CLO	OVER APHID (Therioaphis trifolii)		PEA APHID (Acrythosiphon pisum)	
CLOVER SEE	D MIDGE (Dasineura leguminicola)		OTHER (Specify)	
CLOVER LEA	AFHOPPER (Aceratagallia sanguinolen:	tə)		
12. Indicate the variety most of	closely resembling the application	variety for th	e following:	
CHARACTER	VARIETY	СНА	RACTER	VARIETY
LEAFLET SHAPE	Arlington	SEED COI	-OR	KENSTAR
CUTTING RECOVERY	KENSTAR	LATE SEA	ASON GROWTH	ARLINGTON
WINTER HARDINESS	ARLINGTON	PERSISTE	NCE	KENSTAR
·				

REFERENCES:

Hawkins, R. P. 1953. Investigations on local strains of herbage plants II. Types of red clover and their identification. J. Brit. Grassland Soc. 8, 213-218

Williams, R. D. 1927. Red clover investigations, 1919 - 1926. Welsh Plant Breeding Station Bull., Ser. H. No. 7.

COMMENTS: (If additional space is necessary, use reverse side)

RUBY RED CLOVER

EXHIBIT D

ADDITIONAL DESCRIPTION OF THE VARIETY

Ruby is a variety of two-cut, medium red clover. Ruby has shown superior forage yield to Lakeland and has yielded as well or better than Arlington. Also Ruby has more southern anthracnose resistance than Arlington. It is resistant to powdery mildew and has more resistance to northern anthracnose than Redland or Kenstar. It has good winter survival and persists in the field as well as Arlington. Becuase it has both southern and northern anthracnose resistance, Ruby has a much larger geographic adaptation than either Kenstar or Arlington.

EXHIBIT 1

RED CLOVER VARIETY TRIAL:

University of Wisconsin Trial

Arlington, Wisconsin

Planted: May 1975*

4 reps, 3 cuts

	1975	1976 Yield			
	Powdery	tons/acre	% Stand	% Stand	6-7-76
	Mildew	Dry matter	4-26-76	10-18-76	Anthracnose
Clovage	2.2	4.52	75	36	3.2
Ruby	2.0	4.82	89	74	2.8
Arlington	1.4	4.64	90	76	1.6
Lakeland	1.4	4.40	74	49 .	. 1.9
Kenstar	4.0	4.46	85	74	4.1
Redland	3.0	4.34	84	68	3.9
Pennscott	3.8	3.86	82	36	4.6
Redman	2.5	4.44	60	25	2.2
Dollard	1.5	4.26	82	· 49	1.5
PSC-1	4.2	4.14	72	38	1.2
Common	4.0	4.23	79	21	3.4
K4-184	1.8	4.09	81	75	3.6
HC 1	'-	4.42	89	78	3.0
HC 4	<u>-</u>	4.61	85	86	1.2
HC 6	<u></u>	4.40	82	82	1.4
LSD (.05)	•5	.32	8	16	. 4
MEAN	2.6	4.37	81	58	2.6
CA	12.0%	5.2%	6.6%	19.6%	11.5%

Design: Randomized complete block with four replications.

Plot Size: 5' x 25'. Seeding method: Broadcast.

Soil Type: Paar silt loam.

Disease ratings: Scale of 1 to 5. 1 - no disease; 5 = severe symptoms.

*No 1975 harvest data due to extremely dry weather. Powdery mildew data only.

RED CLOVER VARIETY TRIAL: Dairyland Seed Co., Inc. - Clinton, Wisconsin

3 cuts/year - Seeded: April 1975

	1975	1976	' 75 -' 76
	t/a	t/a	t/a
Variety	D.M.	D.M.	D.M.
Clovage	2.39	3.88	3.14
Ruby	2.73	4.02	3.38
Arlington	2.59	3.88	3.24
Lakeland	2.53	3.81	3.17
EXP-MLM	1.75	3.84	2,79
LSD (.10)	.12	.20	.14
LSD (.05)	.16	.26	.19
MEAN	2.40	3.89	3.14

Design: Randomized complete block with four replications.

Plot Size: 5' x 25'. Seeding method: Broadcast.

Soil type: Miami silt loam.

TABLE 15. RED CLOVER VARIETY TRIAL, EXPERIMENT 338-046
Agronomy South Farm, Urbana, Illinois
1976 Yield Data - Seeded April 11, 1975

	D.M. Yield					
Variety	5/26	7/7	8/4	9/8	Total 1976	Average 2 Yr.
	T/A	T/A	T/A	T/A	T/A	T/A
Arlington	2.81	1.59	0.69	0.84	5.93	3.78
Clovage	2.40	1.39	0.59	0.77	5.16	3.51
Ruby*	2.77	1.66	0.77	0.96	6.16	3.95
Florie (K8-113)	2.62	1.63	0.70	1.12	6.07	4.03
Florex (K8-115)	2.78	1.59	0.63	0.75	5.75	3.63
Kenland	2.46	1.58	0.82	1.01	5.88	3.90
Kenstar	2.62	1.65	0.74	1.16	6.17	3.95
Lakeland	2.72	1.47	0.55	0.72	5.46	3.53
Redland	2.76	1.67	0.70	1.06	6.20	4.00
Redland	2.53	1.62	0.67	1.19	6.01	3.92
Redman	2.58	1.74	0.62	1.02	5.96	3.84
E-684	2.60	1.65	0.68	1.06	5.99	3.90
E-685	2.50	1.71	0.72	1.09	6.01	3.92
E-688	2.53	1.67	0.78	0.96	5.94	3.86
L.S.D. 5%	0.24	0.14	0.12	0.19	0.50	0.24
1.%	0.32	0.18	0.17	0.26	0.66	0.32
C.V. %	6.48	5.94	12.64	13.83	5.90	4.38
Mean	2.62	1.62	0.69	0.98	5.92	3.84

Design: Randomized complete block with four replications.

Plot Size: 5' x 25'. Seeding method: Broadcast. Soil type: Flanagan silt loam and Sidell silt loam.

Fertilization: 0-66-300 after first cutting.

^{*}Ruby was originally listed as the experimental Joy.